

Triviality of Higher Derivative Theories

Victor O. Rivelles*

*Center for Theoretical Physics
Massachusetts Institute of Technology
Cambridge, MA 02139, USA*

We show that some higher derivative theories have a BRST symmetry. This symmetry is due to the higher derivative structure and is not associated to any gauge invariance. If physical states are defined as those in the BRST cohomology then the only physical state is the vacuum. All negative norm states, characteristic of higher derivative theories, are removed from the physical sector. As a consequence, unitarity is recovered but the S-matrix is trivial. We show that a class of higher derivative quantum gravity theories have this BRST symmetry so that they are consistent as quantum field theories. Furthermore, this BRST symmetry may be present in both relativistic and non-relativistic systems.

PACS numbers: 11.10.-z, 11.15.-q, 11.30.Ly, 04.60.-m

Higher derivative (HD) theories were introduced in quantum field theory in an attempt to get rid of ultra-violet divergences [1]. It was soon recognized that they have an energy which is not bounded from below [2] and that these negative energy states can be traded by negative norm states (or ghosts) [3] leading to non unitary theories. Although inconsistent, HD theories have better renormalization properties than conventional ones and a large amount of work was dedicated to the study of such theories [4]. In the gravitational context, for instance, it is able to produce a renormalizable quantum gravity theory [5]. To overcome the ghost problem many attempts were done mainly by imposing some superselection rule or some subsidiary condition to remove the undesirable states [4]. However, there is no general scheme to remove the ghosts and this is done in a case by case basis.

In this letter we will show that some HD theories have a BRST symmetry after ghost fields are introduced. Usually the BRST symmetry is found in gauge theories as a symmetry of the gauge fixed action [6]. Its purpose is to remove the negative norm states associated to the gauge invariance. Physical states are then defined as those which have zero ghost number and are invariant under the BRST symmetry. In the present case, however, the BRST symmetry is not due to gauge invariance. Rather, it is a feature of the HD structure of the action. As it will be shown, it can be found even in the case of a HD real scalar field. Since we have a BRST symmetry it is natural to require that the physical states are those which have zero ghost number and are left invariant by this symmetry, in analogy to what is done in gauge theories. With these requirements we find that HD theories have only one physical state, the vacuum, since all others physical states appear in zero norm combinations. Therefore, the resulting theory is ghost free but it is also empty. In

this way the unitarity problem associated to HD theories can be overcome but the resulting theory has no states besides the vacuum. We will show that this symmetry is present in HD non-Abelian gauge theories and also in HD gravity theories, so that they are unitary but have a trivial S-matrix.

Even though we work mostly with relativistic systems, relativistic invariance is not a necessary ingredient. The HD BRST symmetry may also be found in non-relativistic situations. We will exemplify this for the case of the HD harmonic oscillator.

Let us first consider a real scalar field ϕ in d dimensions with an action

$$S = \int d^d x \left(\frac{1}{2} \mathcal{O}^n \phi \mathcal{O}^n \phi - \bar{c} \mathcal{O}^n c \right), \quad (1)$$

where $\mathcal{O} = \prod_{i=1}^N (\square + m_i^2)$ is a product of N Klein-Gordon operators with masses m_i , c is a ghost, $\bar{c}$ an anti-ghost and n is an integer. This HD action is invariant under the following BRST transformations

$$\delta \phi = c, \quad \delta c = 0, \quad \delta \bar{c} = \mathcal{O}^n \phi. \quad (2)$$

These transformations are nilpotent for ϕ and c and on-shell nilpotent for $\bar{c}$. To find out the role of this symmetry let us introduce an auxiliary field b and rewrite the action (1) as

$$S = \int d^d x \left(b \mathcal{O}^n \phi - \frac{1}{2} b^2 - \bar{c} \mathcal{O}^n c \right). \quad (3)$$

The BRST transformations now read

$$\delta \phi = c, \quad \delta c = 0, \quad \delta \bar{c} = b, \quad \delta b = 0, \quad (4)$$

and are off-shell nilpotent. This action can now be written as a total BRST transformation

$$S = \int d^d x \delta \left[\bar{c} \left(\mathcal{O}^n \phi - \frac{1}{2} b \right) \right], \quad (5)$$

indicating that theory may be empty. It looks like a topological theory of Witten type [7] but it clearly depends on

*On leave from Instituto de Física, Universidade de São Paulo, Caixa Postal 66318, 05315-970, São Paulo, SP, Brazil; Electronic address: rivelles@lns.mit.edu

the local structure of the space-time. In topological theories there are no local degrees of freedom from the start while here we have positive and negative norm states before introducing the ghost fields. The requirement that the physical states are in the cohomology of the BRST charge means that all of them, except the vacuum, appear in zero norm combinations as we will show.

Interactions can be introduced through a prepotential $U(\phi)$. The action

$$S = \int d^d x \delta \left[\bar{c}(\mathcal{O}^n \phi - \frac{1}{2}b - U) \right], \quad (6)$$

which is invariant under the BRST transformations (4), yields, after elimination of the auxiliary field,

$$S = \int d^d x \left[\frac{1}{2}(\mathcal{O}^n \phi)^2 + \frac{1}{2}U^2 + U\mathcal{O}^n \phi - \bar{c}(\mathcal{O}^n c + U'c) \right]. \quad (7)$$

Assuming that the BRST symmetry has no anomaly it is enough to examine the free case to find out the physical states.

In order to simplify the analysis we will consider from now on the case $n = 1$ and only one factor in $\mathcal{O}$, that is $N = 1$. The simplest situation is the one dimensional case, that is, quantum mechanics, where $\mathcal{O} = d^2/dt^2 + m^2$. Then the action (1) reduces to that of a HD harmonic oscillator. Canonical quantization is straightforward if we keep the auxiliary field and use the action in the form (3) where only first order derivatives appear after an integration by parts is performed. We then find that the non trivial equal time commutation relations are

$$[\phi, \dot{b}] = [b, \dot{\phi}] = -i, \quad (8)$$

where dots denote time derivatives. The solution to the field equation for ϕ and b can be expanded as

$$\phi(t) = \frac{1}{2m^{3/2}} [(a + bmt)e^{-imt} + (a^\dagger + b^\dagger mt)e^{imt}], \quad (9)$$

$$b(t) = -im^{1/2} (be^{-imt} - b^\dagger e^{imt}). \quad (10)$$

The canonical Hamiltonian yields

$$H = m(2b^\dagger b - ib^\dagger a + ia^\dagger b), \quad (11)$$

and using the transformation $\tilde{a} = ia - b/2$, $\tilde{b} = ia - 3b/2$ we find

$$H = m(\tilde{a}^\dagger \tilde{a} - \tilde{b}^\dagger \tilde{b}), \quad (12)$$

so that it is not positive definite at the classical level. Upon quantization we find that the non-trivial commutators for the operators a and b are given by

$$[a, a^\dagger] = -1, \quad [a, b^\dagger] = i, \quad (13)$$

where a and b are annihilation operators and $a^\dagger$ and $b^\dagger$ are creation ones. By the transformation $\hat{a} = b - ia$, $\hat{b} = -ia$, we find

$$[\hat{a}, \hat{a}^\dagger] = -[\hat{b}, \hat{b}^\dagger] = 1, \quad (14)$$

so that the Hilbert space is not positive definite.

We now consider the ghost fields in (1). Quantization is straightforward. The solution to the equations of motion can be expanded as

$$c(t) = \frac{1}{2m^{3/2}} (ce^{-i\omega t} + c^\dagger e^{i\omega t}), \quad (15)$$

$$\bar{c}(t) = m^{1/2} (\bar{c}e^{-i\omega t} + \bar{c}^\dagger e^{i\omega t}), \quad (16)$$

where c and $\bar{c}$ are annihilation operators and $c^\dagger$ and $\bar{c}^\dagger$ are creation operators. The non trivial anti-commutators are given by

$$\{c, \bar{c}^\dagger\} = -\{\bar{c}, c^\dagger\} = 1. \quad (17)$$

The BRST transformations (4) for the operators in (9), (10), (15) and (16) (not for the fields) are

$$\delta a = c, \quad \delta c = 0, \quad \delta \bar{c} = -ib, \quad \delta b = 0. \quad (18)$$

This means that these operators belong to the quartet representation of the BRST algebra [8]. As a consequence, the BRST invariant states, build out of these operators, appear only in zero norm combinations through the quartet mechanism. Therefore the only physical state is the vacuum. In this way the HD harmonic oscillator is free of negative norm states but the only state left is the vacuum.

The situation for the scalar field is analogous to the quantum mechanical case. Now $\mathcal{O} = \square + m^2$ and the fields have expansions similar to (9), (10), (15) and (16) with the creation and annihilation operators depending on the momenta. They obey commutation relations similar to (13) and (17) and the Hamiltonian has the same form as (11). Hence the BRST transformations are the same as in (18) and the quartet mechanism is also operational in this case. The only physical state is the vacuum. The massless case needs some care but again the same result is found.

At this point it is worth to remind that to remove the negative norm states for the HD scalar field theory the condition $b = 0$ is often employed [9]. It is argued that there is a “gauge symmetry” $\delta\phi = \Lambda$ with Λ satisfying $(\square + m^2)\Lambda = 0$. The “gauge condition” $b = 0$ is then imposed to remove the ghost states. However, the Hamiltonian analysis reveals that there is no true gauge symmetry since there are no constraints at all. Amazingly, a “BRST symmetry” associated to this “gauge symmetry” can be found [10] and it agrees with (18). Now its origin is clear; it is completely due to the HD structure as we have shown.

We can now easily generalize this construction to other types of fields. Since we want to keep the same BRST transformations (4) then the ghosts and the auxiliary field b must have the same tensor structure as the field under consideration. For gauge theories we should also take into account the ordinary Faddeev-Popov ghosts associated to the gauge symmetry since they have a different origin from the ghosts coming from the HD structure

[11]. We will argue, however, that only one set of ghost fields is needed.

For a theory with a non-Abelian vector field we start with

$$S = \int d^d x \text{Tr} \delta \left[\bar{c}^\mu (E_\mu - \frac{1}{2} b_\mu) \right], \quad (19)$$

where E_μ depends only on the vector field A_μ . The BRST transformations are given by

$$\delta A_\mu = c_\mu, \quad \delta c_\mu = 0, \quad \delta \bar{c}_\mu = b_\mu, \quad \delta b_\mu = 0. \quad (20)$$

This structure is similar to that found in topological field theories [12]. There, E, b and $\bar{c}$ are two forms instead of vectors so that a topological invariant is generated after the elimination of the auxiliary field. Here the tensor structure is quite different and E_μ is chosen so that it gives rise to a HD theory. For a fourth order gauge theory we can choose

$$E_\mu = D^\nu F_{\nu\mu} + \frac{1}{2\xi} \partial_\mu \partial^\nu A_\nu. \quad (21)$$

The first term in (21) will produce a gauge invariant term in the action if the second term is absent. The second term breaks gauge invariance and can be regarded as a gauge fixing term since it allows us to find the propagator for A_μ . At this point we should decide whether we introduce the gauge fixing term in E_μ , as in (21), or if we consider the ordinary Faddeev-Popov procedure for gauge fixing. If we do not add the gauge fixing term in (21) then E_μ will give rise to an action for $\bar{c}_\mu$ and c_μ which has a gauge symmetry $\delta \bar{c}_\mu = D_\mu \bar{\Sigma}$, $\delta c_\mu = D_\mu \Sigma$, where $\bar{\Sigma}$ and Σ are Grassmannian functions. This would require the introduction of ghosts for ghosts besides the ordinary Faddeev-Popov ghosts for A_μ . We then choose to introduce the gauge fixing term directly in E_μ to avoid the proliferation of ghost fields.

With the choice (21) we find that (19) yields the following action

$$\begin{aligned} S = & \frac{1}{2} \int d^d x \text{Tr} \left(D^\nu F_{\nu\mu} D_\rho F^{\rho\mu} + \frac{1}{4\xi^2} \partial^\mu \partial A \partial_\mu \partial A \right. \\ & + \frac{1}{\xi} D^\nu F_{\nu\mu} \partial^\mu \partial A + \bar{\mathcal{F}}^{\mu\nu} \mathcal{F}_{\mu\nu} - 2i \{ \bar{c}^\mu, c^\nu \} F_{\mu\nu} \\ & \left. + \frac{1}{\xi} \partial^\mu \bar{c}_\mu \partial^\nu c_\nu \right), \end{aligned} \quad (22)$$

where $\bar{\mathcal{F}}^{\mu\nu}$ and $\mathcal{F}_{\mu\nu}$ are the field strengths for $\bar{c}^\mu$ and c_μ , respectively. Canonical quantization is easily performed keeping the auxiliary field b_μ since it gives rise to a first order Lagrangian. No constraints are found and the Hamiltonian is not positive definite. Quantization simplifies in the Feynman gauge $\xi = 1$ and we find that the quartet mechanism applies to each component of the vector field. Again, the only physical state is the vacuum. Coupling to ordinary or HD matter is straightforward. However, only non-minimal couplings arise. Details will be given elsewhere.

Other choices for E_μ are possible. For instance,

$$E_\mu = \frac{1}{m} D^\nu F_{\nu\mu} + \frac{m}{2} A_\mu, \quad (23)$$

gives rise to a fourth order massive vector theory which includes the standard Maxwell term. However, there is no gauge invariance in this case.

For gravity the ghosts and the auxiliary field are second order symmetric tensors. The action is

$$S = \int d^d x \delta \left[\bar{c}^{\mu\nu} (E_{\mu\nu} - \frac{1}{2} b_{\mu\nu}) \right], \quad (24)$$

and the BRST transformations are given by

$$\delta g_{\mu\nu} = c_{\mu\nu}, \quad \delta c_{\mu\nu} = 0, \quad \delta \bar{c}^{\mu\nu} = b^{\mu\nu}, \quad \delta b^{\mu\nu} = 0. \quad (25)$$

For a generic fourth order gravity theory we can choose

$$E_{\mu\nu} = g^{1/4} (c_1 R_{\mu\nu} + c_2 g_{\mu\nu} R + c_3 g_{\mu\nu} + \dots), \quad (26)$$

where $g_{\mu\nu}$ is the metric, g its determinant, $R_{\mu\nu}$ the Ricci tensor, R the curvature scalar and dots denote the gauge fixing terms. After the elimination of the auxiliary field we find that the action for the gravitational sector can be written as

$$S_{gr} = \int d^d x \sqrt{-g} [\gamma R - \Lambda + \alpha R^2 + \beta R^{\mu\nu} R_{\mu\nu}], \quad (27)$$

with $\gamma^2 = -4\Lambda(d\alpha + \beta)/d$. Gauge fixing terms were omitted. Notice that we can not get the Einstein-Hilbert action since setting $\alpha = \beta = 0$ to get rid of the HD contributions also eliminates the term in R . However, we can choose $\Lambda = 0$ in order to get a purely HD gravity theory. Alternatively we can choose $\beta = -d\alpha$, which corresponds to a traceless Ricci tensor. This case is distinguished by energy considerations [13]. The conformal case has no special features. Notice also that (26) does not allow us to generate the square of the Riemann tensor thus precluding Gauss-Bonnet terms which arise in string theory. Details will be presented elsewhere.

Some final remarks are in order. Could a BRST symmetry be found for ordinary second order theories? Covariance requires that the operator $\mathcal{O}$ be at least quadratic in the derivatives. However, in lower dimensions, this requirement can be relaxed. Chiral bosons in two dimensions have a BRST symmetry which allows the vacuum as the only physical state [14]. In higher dimensions we could consider the action (6) without the term $\mathcal{O}^n \phi$ and U as a function of ϕ and its derivatives. Then we can take $U = \sqrt{\phi(\Box + m^2)\phi}$. The term U^2 in (7) yields the Lagrangian for the ordinary scalar field but we get non-local interactions with the ghosts due to the term $U' \bar{c} c$. Even though there is a BRST symmetry, a detailed analysis shows that on-shell $\bar{c} = b = 0$ so that the quartet mechanism can not be applied. It is imperative to have non-trivial solutions to the field equation for the quartet mechanism to work. Since $\bar{c} = 0$ all contributions from the ghost sector to the physical Hilbert

space vanish because they must be functions of $\bar{c}c$. Then we end up with the usual scalar theory and the BRST symmetry is trivial in this case.

A more radical possibility would be to consider non-local expression for $\mathcal{O}$, for instance $\mathcal{O} = \sqrt{\square + m^2}$. The action (1) with $n = 1$ would be the ordinary one for a scalar field but for the ghosts we find a non-local action. Quantization of such non-local theories has been done [15] but it not clear how the quartet mechanism can be applied with non-local ghosts. This deserves further understanding.

Returning to the quantum mechanical case, the action (7) resembles that of supersymmetric quantum mechanics if $\mathcal{O} = d/dt$ and the ghosts are regarded as fermions. The BRST transformations (4) are identical to the supersymmetry transformations but the anti-BRST transformation (obtained by replacing ghosts by anti-ghosts and vice-versa) $\bar{\delta}\phi = \bar{c}$, $\bar{\delta}\bar{c} = 0$, $\bar{\delta}c = -(\mathcal{O} + V)$, are not. The anti-commutator of the BRST and anti-BRST transformations vanishes as expected while in the supersymmetric case it would be proportional to the Hamiltonian. There are crucial signs in the action and in the transformation rules which, when changed, produces the supersymmetric model. Hence the supersymmetric model is obtained by twisting the BRST symmetry.

We should also remark that the BRST symmetry can not be found in any HD theory. For instance, in the simple case of non-degenerated masses

$$S = \int d^d x \left[\frac{1}{2}(\square + m_1^2)\phi(\square + m_2^2)\phi - \bar{c}(\square + m_1^2)c \right], \quad (28)$$

there is a symmetry of the action given by $\delta\phi = c$, $\delta c = 0$, $\delta\bar{c} = (\square + m_2^2)\phi$. However, this symmetry is not nilpotent since on-shell $\delta^2\bar{c} = (m_2^2 - m_1^2)c \neq 0$. It is nilpotent only when both masses are equal thus implying in (1) with $n = 1$ and $\mathcal{O} = \square + m^2$, where m is the common mass.

As a last remark it must be stressed that a trivial topology for the space (or space-time) was assumed throughout the paper. If a non-trivial topology is present then topological excitations may arise as they do in topological field theories of Witten type. For instance, in the case of non-Abelian gauge fields (22) there are instanton solutions which may give non-trivial contributions to correlation functions. This is presently under investigation and will be reported elsewhere.

Acknowledgments

I would like to thank A. J. Accioly, S. Deser, R. Jackiw, I. Shapiro and S. Sorella for useful discussions. This work was partially supported by CAPES, CNPQ and PRONEX under contract CNPq 66.2002/1998-99.

-
- [1] W. Thirring, Phys. Rev. **77**, 570 (1950).
 - [2] A. Pais and G. E. Uhlenbeck, Phys. Rev. **79**, 145 (1950).
 - [3] W. Heisenberg, Nucl. Phys. **4**, 532 (1957).
 - [4] See the following papers for earlier references: J. Lukierski, Acta Phys. Polon. **32**, 771 (1967); N. Nakanishi, Prog. Theor. Phys. Suppl. **51**, 1 (1972); K. S. Stelle, Gen. Rel. Grav. **9**, 353 (1978); H. Narnhofer and W. Thirring, Phys. Lett. **76B**, 428 (1978); M. Mintchev, J. Phys. A **13** (1980) 1841; E. S. Fradkin and A. A. Tseytlin, Nucl. Phys. B **201**, 469 (1982); P. Broadbridge, J. Phys. A **16**, 3271 (1983); N. H. Barth and S. M. Christensen, Phys. Rev. D **28**, 1876 (1983); U. Moschella, J. Math. Phys. **31**, 2480 (1990); I. L. Buchbinder, S. D. Odintsov and I. L. Shapiro, "Effective Action In Quantum Gravity," (IOP, Bristol, 1992); M. Asorey, J. L. Lopez and I. L. Shapiro, Int. J. Mod. Phys. A **12**, 5711 (1997) [arXiv:hep-th/9610006]; F. J. de Urries, J. Julve and E. J. Sanchez, J. Phys. A **34**, 8919 (2001) [arXiv:hep-th/0105301]; S. W. Hawking and T. Hertog, Phys. Rev. D **65**, 103515 (2002) [arXiv:hep-th/0107088]; T. C. Cheng, P. M. Ho and M. C. Yeh, Nucl. Phys. B **625**, 151 (2002) [arXiv:hep-th/0111160]; A. Accioly, A. Azeredo and H. Mukai, J. Math. Phys. **43**, 473 (2002); E. J. Villasenor, J. Phys. A **35**, 6169 (2002) [arXiv:hep-th/0203197]; S. De Filippo and F. Maimone, AIP Conf. Proc. **643**, 373 (2003) [arXiv:gr-qc/0207059]; G. Magnano and L. M. Sokolowski, arXiv:gr-qc/0209022; S. Nojiri, arXiv:hep-th/0210056.
 - [5] K. S. Stelle, Phys. Rev. D **16**, 953 (1977).
 - [6] The BRST transformations may be written in different ways (even in non local form) but that is not the case here. This is discussed in V. O. Rivelles, Phys. Rev. Lett. **75**, 4150 (1995) [arXiv:hep-th/9509028]; V. O. Rivelles, Phys. Rev. D **53**, 3247 (1996) [arXiv:hep-th/9510136]; V. O. Rivelles, Class. Quant. Grav. **19**, 2525 (2002) [arXiv:hep-th/0109171].
 - [7] D. Birmingham, M. Blau, M. Rakowski and G. Thompson, Phys. Rept. **209**, 129 (1991).
 - [8] T. Kugo and I. Ojima, Prog. Theor. Phys. Suppl. **66**, 1 (1979).
 - [9] D. Zwanziger, Phys. Rev. D **17**, 457 (1978); A. Z. Capri, G. Grubl and R. Kobes, Annals Phys. **147**, 140 (1983); U. Moschella and F. Strocchi, Lett. Math. Phys. **19** (1990) 143; U. Moschella and F. Strocchi, Lett. Math. Phys. **19** (1990) 143.
 - [10] In fact, it was found in the singleton field theory in M. Flato and C. Fronsdal, Phys. Lett. B **189**, 145 (1987).
 - [11] A. Gavrielides, T. K. Kuo and S. Y. Lee, Phys. Rev. D **13**, 2912 (1976).
 - [12] R. Brooks, D. Montano and J. Sonnenschein, Phys. Lett.

- B **214**, 91 (1988).
- [13] S. Deser and B. Tekin, Phys. Rev. Lett. **89**, 101101 (2002) [arXiv:hep-th/0205318]; S. Deser and B. Tekin, arXiv:hep-th/0212292.
- [14] H. O. Girotti, M. Gomes and V. O. Rivelles, Phys. Rev. D **45**, 3329 (1992) [arXiv:hep-th/9202043].
- [15] D. G. Barci, L. E. Oxman and M. Rocca, Int. J. Mod. Phys. A **11**, 2111 (1996) [arXiv:hep-th/9503101]; D. G. Barci and L. E. Oxman, Mod. Phys. Lett. A **12**, 493 (1997) [arXiv:hep-th/9611147].